

Title	Some remarks on the relative polar variety and the Brasselet number (Research on topology and differential geometry using singularity theory of differentiable maps)
Author(s)	de Carvalho Santana, Hellen Monção
Citation	数理解析研究所講究録 = RIMS Kokyuroku (2019), 2140: 58-70
Issue Date	2019-12
URL	<a href="http://hdl.handle.net/2433/254923">http://hdl.handle.net/2433/254923</a>
Right	
Type	Departmental Bulletin Paper
Textversion	publisher

# Some remarks on the relative polar variety and the Brasselet number \*

Hellen Santana

## Introduction

Let  $f : (\mathbb{C}^n, 0) \rightarrow (\mathbb{C}, 0)$  be an analytic function defined in a neighborhood of the origin and  $\Sigma f$  the critical locus of  $f$ . The Milnor fiber  $F_{f,0}$  is given by  $f^{-1}(\delta) \cap B_\epsilon$ , where  $\delta$  is a regular value of  $f$ ,  $0 < |\delta| \ll \epsilon \ll 1$ . In [15], Milnor proved that, if  $f$  has an isolated singularity,  $F_{f,0}$  has the homotopy type of a wedge of  $\mu(f)$  spheres of dimension  $n - 1$ , where  $\mu(f)$  is the Milnor number of  $f$ . Also,  $\mu(f)$  is the number of Morse points in a Morsefication of  $f$  in a neighborhood of the origin.

In [6], Hamm generalized Milnor's results for complete intersections with isolated singularity  $F = (f_1, \dots, f_k) : (\mathbb{C}^n, 0) \rightarrow (\mathbb{C}^k, 0)$ ,  $1 < k < n$ , proving that the Milnor fiber  $F^{-1}(\delta) \cap B_\epsilon$ ,  $0 < |\delta| \ll \epsilon \ll 1$ , has the homotopy type of a wedge of  $\mu(F)$  spheres of dimension  $n - k$ . In this context, Lê [7] and Greuel [5] proved that  $\mu(F) + \mu(F') = \dim_{\mathbb{C}} \left( \frac{\mathcal{O}_{\mathbb{C}^n, 0}}{I} \right)$ , where  $F' : (\mathbb{C}^n, 0) \rightarrow (\mathbb{C}^{k-1}, 0)$  is the map with components  $f_1, \dots, f_{k-1}$  and  $I$  is the ideal generated by  $f_1, \dots, f_{k-1}$  and the  $(k \times k)$ -minors  $\frac{\partial(f_1, \dots, f_k)}{\partial(x_{i_1}, \dots, x_{i_k})}$ . Notice that the number  $\dim_{\mathbb{C}} \left( \frac{\mathcal{O}_{\mathbb{C}^n, 0}}{I} \right)$  is the number of critical points of a Morsefication of  $f_k$  appearing on the Milnor fibre of  $F'$ .

If  $f$  is defined over a complex analytic space  $X$  and  $f$  has an isolated singularity at the origin, a generalization for the Milnor number is the Euler obstruction of the function  $f$ , introduced in [2], by Brasselet, Massey, Parameswaran and Seade. In [17], Seade, Tibăr and Verjovsky proved that, up to sign, this number is the number of Morse critical points of a stratified Morsefication of  $f$  appearing in the regular part of  $X$ .

In a more general context, if  $f$  is defined over a complex analytic space  $X$  equipped with a good stratification  $\mathcal{V}$  of  $X$  relative to  $f$  (see Definition 2.1) and the function  $f$  does not have isolated singularity at the origin, a way to describe the generalized Milnor fiber  $X \cap f^{-1}(\delta) \cap B_\epsilon$  is to use the Brasselet number of  $f$  at the origin,  $B_{f,X}(0)$ , introduced by Dutertre and Grulha, in [3], and that generalizes the Milnor number to this more general setting and the local Euler obstruction, introduced by MacPherson, in [10], in his proof for the Deligne-Grothendieck conjecture. In [3], the authors presented several formulas to compute Brasselet numbers counting number of stratified Morse critical points. For example, they present a Lê-Greuel type formula for the Brasselet number: if  $g : X \rightarrow \mathbb{C}$  is prepolare with respect to  $\mathcal{V}$  at the origin (see Definition 2.4) and  $0 < |\delta| \ll \epsilon \ll 1$ , then

---

\*Research partially supported by FAPESP - Brazil, Grants 2015/25191-9 and 2017/18543-1.

Key-words: Brasselet number, relative polar variety, local Euler obstruction, stratified Morse critical points

$$B_{f,X}(0) - B_{f,X^g}(0) = (-1)^{d-1}n_q,$$

where  $n_q$  is the number of Morse critical points of a partial Morsefication of  $g|_{X \cap f^{-1}(\delta) \cap B_\epsilon}$  appearing in the regular part of  $X$ , and  $X^g = X \cap \{g = 0\}$ .

They also proved results about the topology of functions with isolated singularity defined over an analytic complex Whitney stratified variety  $X$ . If  $X$  is equidimensional, let  $f, g : X \rightarrow \mathbb{C}$  be analytic functions with isolated singularity at the origin such that  $g$  is prepolar at the origin with respect to the good stratification induced by  $f$  (see (1)) and  $f$  is prepolar at the origin with respect to the good stratification induced by  $g$ , then

$$B_{f,X}(0) - B_{g,X}(0) = (-1)^{d-1}(n_q - m_q).$$

where  $X^f = X \cap \{f = 0\}$ ,  $n_q$  is the number of Morse critical points of a Morsefication of  $g|_{X \cap f^{-1}(\delta) \cap B_\epsilon}$  appearing in the regular part of  $X$  and  $m_q$  is the number of Morse critical points of a Morsefication of  $f|_{X \cap g^{-1}(\delta) \cap B_\epsilon}$  appearing in the regular part of  $X$ , for  $0 < |\delta| \ll \epsilon \ll 1$ .

Computing these numbers of stratified Morse critical points is directly connected to relative polar varieties. Consider  $l$  a linear form in  $\mathbb{C}^n$ ,  $\mathcal{W}$  a Whitney stratification of an open subset  $U$  of  $X$ ,  $\{W_i \setminus \{l = 0\}, W_i \cap \{l = 0\} \setminus \{0\}, \{0\}\}$  the good stratification of  $U$  induced by  $l$  and a function-germ  $g : (U, 0) \rightarrow (\mathbb{C}, 0)$ . If  $l$  sufficiently generic, the relative polar variety (curve)  $\Gamma_{g,l}$  defined by Lê and Teissier, in [8], coincides with the relative polar curve defined by Massey, in [12], and with the relative polar varieties, defined by Massey, in [14] (see [11] and [13]). Each of these polar varieties are useful not only to compute polar multiplicities ([9]) and intersection numbers ([14]), but also to describe critical loci of pair of functions defined over  $X$  ([12], [3]), which is the approach we are interested the most.

In this work, we use relative polar varieties to compute a number of stratified Morse critical points of a specific type of Morsefication of a function-germ, aiming to obtain informations about the Brasselet number of this germ.

## 1 Local Euler obstruction and Euler obstruction of a function

We begin presenting the local Euler obstruction, a singular invariant defined by MacPherson and used as one of the main tools in his proof for the Deligne-Grothendieck conjecture about the existence and uniqueness of Chern classes for singular varieties.

Let  $(X, 0) \subset (\mathbb{C}^n, 0)$  be an equidimensional reduced complex analytic germ of dimension  $d$  in a open set  $U \subset \mathbb{C}^n$ . Consider a complex analytic Whitney stratification  $\mathcal{V} = \{V_\lambda\}$  of  $U$  adapted to  $X$  such that  $\{0\}$  is a stratum. We choose a small representative of  $(X, 0)$ , denoted by  $X$ , such that  $0$  belongs to the closure of all strata. We write  $X = \bigcup_{i=0}^q V_i$ , where  $V_0 = \{0\}$  and  $V_q = X_{reg}$ , where  $X_{reg}$  is the regular part of  $X$ . We suppose that  $V_0, V_1, \dots, V_{q-1}$  are connected and that the analytic sets  $\overline{V_0}, \overline{V_1}, \dots, \overline{V_q}$  are reduced. We write  $d_i = \dim(V_i)$ ,  $i \in \{1, \dots, q\}$ . Note that  $d_q = d$ .

Let  $G(d, N)$  be the Grassmannian manifold,  $x \in X_{reg}$  and consider the Gauss map  $\phi : X_{reg} \rightarrow U \times G(d, N)$  given by  $x \mapsto (x, T_x(X_{reg}))$ .

**Definition 1.1.** The closure of the image of the Gauss map  $\phi$  in  $U \times G(d, N)$ , denoted by  $\tilde{X}$ , is called **Nash modification** of  $X$ . It is a complex analytic space endowed with an analytic projection map  $\nu : \tilde{X} \rightarrow X$ .

Consider the extension of the tautological bundle  $\mathcal{T}$  over  $U \times G(d, N)$ . Since  $\tilde{X} \subset U \times G(d, N)$ , we consider  $\tilde{T}$  the restriction of  $\mathcal{T}$  to  $\tilde{X}$ , called the **Nash bundle**, and  $\pi : \tilde{T} \rightarrow \tilde{X}$  the projection of this bundle.

In this context, denoting by  $\varphi$  the natural projection of  $U \times G(d, N)$  at  $U$ , we have the following diagram:

$$\begin{array}{ccc} \tilde{T} & \longrightarrow & \mathcal{T} \\ \pi \downarrow & & \downarrow \\ \tilde{X} & \longrightarrow & U \times G(d, N) \\ \nu \downarrow & & \downarrow \varphi \\ X & \longrightarrow & U \subseteq \mathbb{C}^N \end{array}$$

Considering  $\|z\| = \sqrt{z_1 \bar{z}_1 + \dots + z_N \bar{z}_N}$ , the 1-differential form  $w = d\|z\|^2$  over  $\mathbb{C}^N$  defines a section in  $T^*\mathbb{C}^N$  and its pullback  $\varphi^*w$  is a 1-form over  $U \times G(d, N)$ . Denote by  $\tilde{w}$  the restriction of  $\varphi^*w$  over  $\tilde{X}$ , which is a section of the dual bundle  $\tilde{T}^*$ .

Choose  $\epsilon$  small enough for  $\tilde{w}$  be a non zero section over  $\nu^{-1}(z)$ ,  $0 < \|z\| \leq \epsilon$ , let  $B_\epsilon$  be the closed ball with center at the origin with radius  $\epsilon$  and denote:

1.  $Obs(\tilde{T}^*, \tilde{w}) \in \mathbb{H}^{2d}(\nu^{-1}(B_\epsilon), \nu^{-1}(S_\epsilon), \mathbb{Z})$  as the obstruction for extending  $\tilde{T}$  from  $\nu^{-1}(S_\epsilon)$  to  $\nu^{-1}(B_\epsilon)$ ;
2.  $O_{\nu^{-1}(B_\epsilon), \nu^{-1}(S_\epsilon)}$  as the fundamental class in  $\mathbb{H}_{2d}(\nu^{-1}(B_\epsilon), \nu^{-1}(S_\epsilon), \mathbb{Z})$ .

**Definition 1.2.** The **local Euler obstruction** of  $X$  at 0,  $Eu_X(0)$ , is given by the evaluation

$$Eu_X(0) = \langle Obs(\tilde{T}^*, \tilde{w}), O_{\nu^{-1}(B_\epsilon), \nu^{-1}(S_\epsilon)} \rangle.$$

In [1], Brasselet, Lê and Seade proved a formula to make the calculation of the Euler obstruction easier.

**Theorem 1.3.** (Theorem 3.1 of [1]) Let  $(X, 0)$  and  $\mathcal{V}$  be given as before, then for each generic linear form  $l$ , there exists  $\epsilon_0$  such that for any  $\epsilon$  with  $0 < \epsilon < \epsilon_0$  and  $\delta \neq 0$  sufficiently small, the Euler obstruction of  $(X, 0)$  is equal to

$$Eu_X(0) = \sum_{i=1}^q \chi(V_i \cap B_\epsilon \cap l^{-1}(\delta)) \cdot Eu_X(V_i),$$

where  $\chi$  is the Euler characteristic,  $Eu_X(V_i)$  is the Euler obstruction of  $X$  at a point of  $V_i$ ,  $i = 1, \dots, q$  and  $0 < |\delta| \ll \epsilon \ll 1$ .

Let us give the definition of another invariant introduced by Brasselet, Massey, Parameswaran and Seade in [2]. Let  $f : X \rightarrow \mathbb{C}$  be a holomorphic function with isolated singularity at the origin given by the restriction of a holomorphic function  $F : U \rightarrow \mathbb{C}$  and denote by  $\bar{\nabla}F(x)$  the conjugate of the gradient vector field of  $F$  in  $x \in U$ ,

$$\bar{\nabla}F(x) := \left( \frac{\partial \bar{F}}{\partial x_1}, \dots, \frac{\partial \bar{F}}{\partial x_n} \right).$$

Since  $f$  has an isolated singularity at the origin, for all  $x \in X \setminus \{0\}$ , the projection  $\hat{\zeta}_i(x)$  of  $\bar{\nabla}F(x)$  over  $T_x(V_i(x))$  is non-zero, where  $V_i(x)$  is a stratum containing  $x$ . Using this projection, the authors constructed, in [2], a stratified vector field over  $X$ , denoted by  $\bar{\nabla}f(x)$ . Let  $\tilde{\zeta}$  be the lifting of  $\bar{\nabla}f(x)$  as a section of the Nash bundle  $\tilde{T}$  over  $\tilde{X}$ , without singularity over  $\nu^{-1}(X \cap S_\epsilon)$ .

Let  $\mathcal{O}(\tilde{\zeta}) \in \mathbb{H}^{2n}(\nu^{-1}(X \cap B_\epsilon), \nu^{-1}(X \cap S_\epsilon))$  be the obstruction cocycle for extending  $\tilde{\zeta}$  as a non zero section of  $\tilde{T}$  inside  $\nu^{-1}(X \cap B_\epsilon)$ .

**Definition 1.4.** The **local Euler obstruction of the function**  $f$ ,  $Eu_{f,X}(0)$  is the evaluation of  $\mathcal{O}(\tilde{\zeta})$  on the fundamental class  $[\nu^{-1}(X \cap B_\epsilon), \nu^{-1}(X \cap S_\epsilon)]$ .

The next theorem compares the Euler obstruction of a space  $X$  with the Euler obstruction of function defined over  $X$ .

**Theorem 1.5.** (Theorem 3.1 of [2]) Let  $(X, 0)$  and  $\mathcal{V}$  be given as before and let  $f : (X, 0) \rightarrow (\mathbb{C}, 0)$  be a function with an isolated singularity at 0. For  $0 < |\delta| \ll \epsilon \ll 1$ , we have

$$Eu_{f,X}(0) = Eu_X(0) - \sum_{i=1}^q \chi(V_i \cap B_\epsilon \cap f^{-1}(\delta)) \cdot Eu_X(V_i).$$

In [17], Seade, Tibăr and Verjovsky proved that the Euler obstruction of a function  $f$  is also related to the number of Morse critical points of a Morsefication of  $f$ . Before we state their result, let us see the definition of a general point.

**Definition 1.6.** Let  $(X, 0) \subset (U, 0)$  be a germ of complex analytic space in  $\mathbb{C}^n$  equipped with a Whitney stratification and let  $f : (X, 0) \rightarrow (\mathbb{C}, 0)$  be an analytic function, given by the restriction of an analytic function  $F : (U, 0) \rightarrow (\mathbb{C}, 0)$ . Then 0 is said to be a **generic point** of  $f$  if the hyperplane  $\text{Ker}(d_0F)$  is transverse in  $\mathbb{C}^n$  to all limit of tangent spaces  $\lim_{n \rightarrow \infty} T_{x_n}(V_\alpha)$ , for all  $V_\alpha$  and sequence of points  $x_n \in V_\alpha$  converging to 0.

Now, let us see the definition of a Morsefication of a function.

**Definition 1.7.** A function  $f : (X, 0) \rightarrow (\mathbb{C}, 0)$  is said to be **Morse stratified** if for all strata  $V_\alpha$ , with  $\dim V_\alpha \geq 1$ , 0 is a generic point of the restriction  $f|_{V_\alpha}$  and for  $V_0 = \{0\}$ , 0 is a Morse point of  $f|_{V_0}$ .

A stratified Morsefication of a germ of analytic function  $f : (X, 0) \rightarrow (\mathbb{C}, 0)$  is a deformation  $\tilde{f}$  of  $f$  such that  $\tilde{f}$  is Morse stratified.

**Proposition 1.8.** (Proposition 2.3 of [17]) Let  $f : (X, 0) \rightarrow (\mathbb{C}, 0)$  be a germ of analytic function with isolated singularity at the origin. Then,

$$Eu_{f,X}(0) = (-1)^{d_{n_{reg}}},$$

where  $n_{reg}$  is the number of Morse points in  $X_{reg}$  in a stratified Morsefication of  $f$ .

In the case where we have a function with several number of isolated critical points, one can be interested in a deformation of this function which is a Morsefication around each one of these singularities. This is what Dutertre and Grulha called a partial Morsefication.

**Definition 1.9.** A **partial Morsefication** of  $g : f^{-1}(\delta) \cap X \cap B_\epsilon \rightarrow \mathbb{C}$  is a function  $\tilde{g} : f^{-1}(\delta) \cap X \cap B_\epsilon \rightarrow \mathbb{C}$  (not necessarily holomorphic) which is a local Morsefication of all isolated critical points of  $g$  in  $f^{-1}(\delta) \cap X \cap \{g \neq 0\} \cap B_\epsilon$  and which coincides with  $g$  outside a small neighborhood of these critical points.

## 2 Brasselet number

We present now the two most important tools in this work: the relative polar variety, defined by Massey in [12], and the Brasselet number, introduced by Dutertre and Grulha in [3]. We also present formulas proved by Dutertre and Grulha to compute Brasselet numbers by counting numbers of stratified Morse critical points.

Let  $X$  be a reduced complex analytic space (not necessarily equidimensional) of dimension  $d$  in an open set  $U \subseteq \mathbb{C}^n$  and let  $f : (X, 0) \rightarrow (\mathbb{C}, 0)$  be an analytic map. We write  $V(f) = f^{-1}(0)$ .

**Definition 2.1.** A **good stratification of  $X$  relative to  $f$**  is a stratification  $\mathcal{V}$  of  $X$  which is adapted to  $V(f)$  such that  $\{V_\lambda \in \mathcal{V}, V_\lambda \not\subseteq V(f)\}$  is a Whitney stratification of  $X \setminus V(f)$  and such that for any pair  $(V_\lambda, V_\gamma)$  such that  $V_\lambda \not\subseteq V(f)$  and  $V_\gamma \subseteq V(f)$ , the  $(a_f)$ -Thom condition is satisfied, that is, if  $p \in V_\gamma$  and  $p_i \in V_\lambda$  are such that  $p_i \rightarrow p$  and  $T_{p_i}V(f)|_{V_\lambda} - f|_{V_\lambda}(p_i)$  converges to some  $\mathcal{T}$ , then  $T_pV_\gamma \subseteq \mathcal{T}$ .

If  $f : X \rightarrow \mathbb{C}$  has a stratified isolated critical point and  $\mathcal{V}$  is a Whitney stratification of  $X$ , then

$$\{V_\lambda \setminus X^f, V_\lambda \cap X^f \setminus \{0\}, \{0\}, V_\lambda \in \mathcal{V}\} \quad (1)$$

is a good stratification of  $X$  relative to  $f$ , called the good stratification induced by  $f$ .

**Definition 2.2.** The **critical locus of  $f$  relative to  $\mathcal{V}$** ,  $\Sigma_{\mathcal{V}}f$ , is given by the union

$$\Sigma_{\mathcal{V}}f = \bigcup_{V_\lambda \in \mathcal{V}} \Sigma(f|_{V_\lambda}).$$

**Definition 2.3.** If  $\mathcal{V} = \{V_\lambda\}$  is a stratification of  $X$ , the **symmetric relative polar variety of  $f$  and  $g$  with respect to  $\mathcal{V}$** ,  $\tilde{\Gamma}_{f,g}(\mathcal{V})$ , is the union  $\bigcup_\lambda \tilde{\Gamma}_{f,g}(V_\lambda)$ , where  $\Gamma_{f,g}(V_\lambda)$  denotes the closure in  $X$  of the critical locus of  $(f, g)|_{V_\lambda \setminus (X^f \cup X^g)}$ ,  $X^f = X \cap \{f = 0\}$  and  $X^g = X \cap \{g = 0\}$ .

**Definition 2.4.** Let  $\mathcal{V}$  be a good stratification of  $X$  relative to a function  $f : (X, 0) \rightarrow (\mathbb{C}, 0)$ . A function  $g : (X, 0) \rightarrow (\mathbb{C}, 0)$  is **prepolar with respect to  $\mathcal{V}$  at the origin** if the origin is a stratified isolated critical point, that is, 0 is an isolated point of  $\Sigma_{\mathcal{V}}g$ .

**Definition 2.5.** A function  $g : (X, 0) \rightarrow (\mathbb{C}, 0)$  is **tractable at the origin with respect to a good stratification  $\mathcal{V}$  of  $X$  relative to  $f : (X, 0) \rightarrow (\mathbb{C}, 0)$**  if  $\dim_0 \tilde{\Gamma}_{f,g}^1(\mathcal{V}) \leq 1$  and, for all strata  $V_\alpha \subseteq X^f$ ,  $g|_{V_\alpha}$  has no critical points in a neighbourhood of the origin except perhaps at the origin itself.

Another concept useful for this work is the notion of constructible functions. Consider a Whitney stratification  $\mathcal{W} = \{W_1, \dots, W_q\}$  of  $X$  such that each stratum  $W_i$  is connected.

**Definition 2.6.** A constructible function with respect to the stratification  $\mathcal{W}$  of  $X$  is a function  $\beta : X \rightarrow \mathbb{Z}$  which is constant on each stratum  $W_i$ , that is, there exist integers  $t_1, \dots, t_q$ , such that  $\beta = \sum_{i=1}^q t_i \cdot 1_{W_i}$ , where  $1_{W_i}$  is the characteristic function of  $W_i$ .

**Definition 2.7.** The Euler characteristic  $\chi(X, \beta)$  of a constructible function  $\beta : X \rightarrow \mathbb{Z}$  with respect to the stratification  $\mathcal{W}$  of  $X$ , given by  $\beta = \sum_{i=1}^q t_i \cdot 1_{W_i}$ , is defined by  $\chi(X, \beta) = \sum_{i=1}^q t_i \cdot \chi(W_i)$ .

Before we state Dutertre and Grulha results, we need to introduce some definitions about normal Morse data. We cite as main references [4] and [18]. The first concept we present is the complex link, an object analogous to the Milnor fibre, important in the study of complex stratified Morse theory.

Let  $V$  be a stratum of the stratification  $\mathcal{V}$  of  $X$  and let  $x$  be a point of  $V$ . Let  $g : (\mathbb{C}^n, 0) \rightarrow (\mathbb{C}, 0)$  be an analytic complex function-germ such that the differential form  $Dg(x)$  is not a degenerate covector of  $\mathcal{V}$  at  $x$ . Let  $N$  be a normal slice to  $V$  at  $x$ , that is,  $N$  is a closed complex submanifold of  $\mathbb{C}^n$  which is transversal to  $V$  at  $x$  and  $N \cap V = \{x\}$ .

**Definition 2.8.** Let  $B_\epsilon$  be the closed ball of radius  $\epsilon$  centered at  $x$ . The **complex link**  $l_V$  of  $V$  is defined by  $l_V = X \cap N \cap B_\epsilon \cap \{g = \delta\}$ , where  $0 < |\delta| \ll \epsilon \ll 1$ .

The **normal Morse datum**  $NMD(V)$  of  $V$  is the pair of spaces

$$NMD(V) = (X \cap N \cap B_\epsilon, X \cap N \cap B_\epsilon \cap \{g = \delta\}).$$

In Part II, section 2.3 of [4], the authors explained why this two notions are independent of all choices made.

**Definition 2.9.** Let  $\beta : X \rightarrow \mathbb{Z}$  be a constructible function with respect to the stratification  $\mathcal{V}$ . Its normal Morse index  $\eta(V, \beta)$  along  $V$  is defined by

$$\eta(V, \beta) = \chi(NMD(V), \beta) = \chi(X \cap N \cap B_\epsilon, \beta) - \chi(l_V, \beta).$$

In the case where the constructible function is the local Euler obstruction, the following identities are valid ([18], page 34):

$$\eta(V', Eu_{\overline{V}}) = 1, \text{ if } V' = V \text{ and } \eta(V', Eu_{\overline{V}}) = 0, \text{ if } V' \neq V.$$

We present now the definition of the Brasselet number and the main theorems of [3], used as inspiration for this work.

Let  $f : (X, 0) \rightarrow (\mathbb{C}, 0)$  be a complex analytic function germ and let  $\mathcal{V}$  be a good stratification of  $X$  relative to  $f$ . We denote by  $V_1, \dots, V_q$  the strata of  $\mathcal{V}$  that are not contained in  $\{f = 0\}$  and we assume that  $V_1, \dots, V_{q-1}$  are connected and that  $V_q = X_{reg} \setminus \{f = 0\}$ . Note that  $V_q$  could be not connected.

**Definition 2.10.** Suppose that  $X$  is equidimensional. Let  $\mathcal{V}$  be a good stratification of  $X$  relative to  $f$ . The **Brasselet number** of  $f$  at the origin,  $B_{f,X}(0)$ , is defined by

$$B_{f,X}(0) = \sum_{i=1}^q \chi(V_i \cap f^{-1}(\delta) \cap B_\epsilon) Eu_X(V_i),$$

where  $0 < |\delta| \ll \epsilon \ll 1$ .

**Remark:** If  $V_q^i$  is a connected component of  $V_q$ ,  $Eu_X(V_q^i) = 1$ .

Notice that if  $f$  has a stratified isolated singularity at the origin, then  $B_{f,X}(0) = Eu_X(0) - Eu_{f,X}(0)$  (see Theorem 1.5).

In [3], Dutertre and Grulha proved interesting formulas describing the topological relation between the Brasselet number and a number of certain critical points of a special type of deformation of functions. Let us now present some of these results.

Let  $g : (X, 0) \rightarrow (\mathbb{C}, 0)$  be a complex analytic function which is tractable at the origin with respect to  $\mathcal{V}$  relative to  $f$ . Then  $\Gamma_{f,g}$  is a complex analytic curve and for  $0 < |\delta| \ll 1$  the critical points of  $g|_{f^{-1}(\delta) \cap X}$  in  $B_\epsilon$  lying outside  $\{g = 0\}$  are isolated. Let  $\tilde{g}$  be a partial Morsefication of  $g : f^{-1}(\delta) \cap X \cap B_\epsilon \rightarrow \mathbb{C}$  and, for each  $i \in \{1, \dots, q\}$ , let  $n_i$  be the number of stratified Morse critical points of  $\tilde{g}$  appearing on  $V_i \cap f^{-1}(\delta) \cap \{g \neq 0\} \cap B_\epsilon$ .



**Theorem 2.11.** (Theorem 4.2 of [3]) Let  $\beta : X \rightarrow \mathbb{Z}$  be a constructible function with respect to the stratification  $\mathcal{V}$ . Suppose that  $g : (X, 0) \rightarrow (\mathbb{C}, 0)$  is a complex analytic function tractable at the origin with respect to  $\mathcal{V}$  relative to  $f$ . For  $0 < |\delta| \ll \epsilon \ll 1$ , we have

$$\chi(X \cap f^{-1}(\delta) \cap B_\epsilon, \beta) - \chi(X \cap g^{-1}(0) \cap f^{-1}(\delta) \cap B_\epsilon, \beta) = \sum_{i=1}^q (-1)^{d_i-1} n_i \eta(V_i, \beta).$$

In the case that  $\beta = Eu_X$ , the last theorem implies the following.

**Corollary 2.12.** (Corollary 4.3 of [3]) Suppose that  $X$  is equidimensional and that  $g$  is tractable at the origin with respect to  $\mathcal{V}$  relative to  $f$ . For  $0 < |\delta| \ll \epsilon \ll 1$ , we have

$$\chi(X \cap f^{-1}(\delta) \cap B_\epsilon, Eu_X) - \chi(X \cap g^{-1}(0) \cap f^{-1}(\delta) \cap B_\epsilon, Eu_X) = (-1)^{d-1} n_q.$$

If one supposes, in addition, that  $g$  is prepolar, a consequence of this result is a Lê-Greuel type formula for the Brasselet number.

**Theorem 2.13.** (Theorem 4.4 of [3]) Suppose that  $X$  is equidimensional and that  $g$  is prepolar with respect to  $\mathcal{V}$  at the origin. For  $0 < |\delta| \ll \epsilon \ll 1$ , we have

$$B_{f,X}(0) - B_{f,X^g}(0) = (-1)^{d-1} n_q,$$

where  $n_q$  is the number of stratified Morse critical points on the top stratum  $V_q \cap f^{-1}(\delta) \cap B_\epsilon$  appearing in a Morsefication of  $g : X \cap f^{-1}(\delta) \cap B_\epsilon \rightarrow \mathbb{C}$ .

In [3], the authors also related the topology of the generalized Minor fibres of  $f$  and  $g$  and some number of Morse points.

**Theorem 2.14.** Suppose that  $g$  (resp.  $f$ ) is prepolar with respect to the good stratification induced by  $f$  (resp.  $g$ ) at the origin. Let  $\beta : X \rightarrow \mathbb{Z}$  be a constructible function with respect to the Whitney stratification  $\mathcal{V}$ . For  $0 < |\delta| \ll \epsilon \ll 1$ ,

$$\chi(X \cap f^{-1}(\delta) \cap B_\epsilon, \beta) - \chi(X \cap g^{-1}(\delta) \cap B_\epsilon, \beta) = \sum_{i=1}^q (-1)^{d_i-1} (n_i - m_i) \eta(V_i, \beta),$$

where  $n_i$  (resp.  $m_i$ ) is the number of stratified Morse critical points on the stratum  $V_i \cap f^{-1}(\delta) \cap B_\epsilon$  (resp.  $V_i \cap g^{-1}(\delta) \cap B_\epsilon$ ) appearing in a Morsefication of  $g : X \cap f^{-1}(\delta) \cap B_\epsilon \rightarrow \mathbb{C}$  (resp.  $f : X \cap g^{-1}(\delta) \cap B_\epsilon \rightarrow \mathbb{C}$ ).

In the case where  $\beta = Eu_X$ , the last theorem implies the following result.

**Corollary 2.15.** Suppose that  $X$  is equidimensional and that  $g$  (resp.  $f$ ) is prepolar with respect to the good stratification induced by  $f$  (resp.  $g$ ) at the origin. Then

$$B_{f,X}(0) - B_{g,X}(0) = (-1)^{d-1} (n_q - m_q),$$

where  $n_q$  (resp.  $m_q$ ) is the number of stratified Morse critical points on the top stratum  $V_q \cap f^{-1}(\delta) \cap B_\epsilon$  (resp.  $V_q \cap g^{-1}(\delta) \cap B_\epsilon$ ) appearing in a Morsefication of  $g : X \cap f^{-1}(\delta) \cap B_\epsilon \rightarrow \mathbb{C}$  (resp.  $f : X \cap g^{-1}(\delta) \cap B_\epsilon \rightarrow \mathbb{C}$ ).



### 3 Brasselet numbers and empty relative polar varieties

In this final section, we present relations between Brasselet numbers of two function-germs in the case where the relative polar variety associated to these germs is empty.

Let  $(X, 0)$  be a complex analytic space with ambient space  $U \subset \mathbb{C}^n$ . Let  $f, g : (X, 0) \rightarrow (\mathbb{C}, 0)$  be germs of holomorphic functions and let  $\mathcal{V}$  be a good stratification of  $U$  relative to  $f$ . Suppose that  $\Sigma_{\mathcal{V}}g \cap \{f = 0\} = \{0\}$ . We aim to obtain information about the Brasselet number of  $f$  and the Brasselet number of  $g$  in the case where the relative polar variety  $\Gamma_{f,g}(\mathcal{V})$  is empty.

We begin with a description of two relevant subsets of the relative polar variety  $\Gamma_{f,g}(\mathcal{V})$ .

**Proposition 3.1.** *The stratified critical set  $\Sigma_{\mathcal{V}}g$  of  $g$  and the symmetric relative polar variety  $\tilde{\Gamma}_{f,g}(\mathcal{V})$  are subsets of  $\Gamma_{f,g}(\mathcal{V})$ .*

**Proof.** If  $x \in \Sigma_{\mathcal{V}}g$ ,  $d_x \tilde{g}|_{V_\alpha} = 0$ , for a stratum  $V_\alpha \in \mathcal{V}$  containing  $x$  and an analytic extension  $\tilde{g}$  of  $g$  in a neighborhood of  $x$ . If  $V_\alpha \subset \{f = 0\}$ , then  $x = 0$ , since  $\Sigma_{\mathcal{V}}g \cap \{f = 0\} = \{0\}$ . If  $V_\alpha \subset X \setminus \{f = 0\}$ ,

$$rk(d_x \tilde{f}|_{V_\alpha}, d_x \tilde{g}|_{V_\alpha}) \leq 1,$$

where  $\tilde{f}$  is an analytic extension of  $f$  in a neighborhood of  $x$ , that is  $x \in \Sigma(f, g)|_{V_\alpha} = \Sigma(f, g)|_{V_\alpha \setminus \{f=0\}}$ . Therefore,  $x \in \Gamma_{f,g}(V_\alpha)$ .

Furthermore,  $\tilde{\Gamma}_{f,g}(V_\alpha)$  is given by components of  $\Gamma_{f,g}(V_\alpha)$  not contained in  $\{g = 0\}$ , that is,  $\tilde{\Gamma}_{f,g}(V_\alpha) = \overline{\Gamma_{f,g}(V_\alpha) \setminus \{g = 0\}} \subseteq \Gamma_{f,g}(V_\alpha)$ . Therefore,

$$\tilde{\Gamma}_{f,g}(\mathcal{V}) \cup \Sigma_{\mathcal{V}}g \subseteq \Gamma_{f,g}(\mathcal{V}).$$

□

Using this proposition, we obtain the following useful information about the behavior of  $g$  with respect to the stratification  $\mathcal{V}$ .

**Corollary 3.2.** *If  $\Gamma_{f,g}(\mathcal{V})$  is empty, then  $g$  is prepolar at the origin with respect to the good stratification  $\mathcal{V}$  of  $X$  relative by  $f$ .*

**Proof.** By Proposition 3.1, if  $\Gamma_{f,g}(\mathcal{V})$  is empty,  $\Sigma_{\mathcal{V}}g$  is empty, that is,  $g$  has no stratified critical point with respect to  $\mathcal{V}$ . □

Let  $n_i$  be the number of stratified Morse critical points of a Morsefication of  $g : X \cap f^{-1}(\delta) \cap B_\epsilon \rightarrow \mathbb{C}$  in  $V_i \cap f^{-1}(\delta) \cap \{g \neq 0\} \cap B_\epsilon$ , for each  $i \in \{1, \dots, q\}$ . The next proposition uses the relative polar variety  $\Gamma_{f,g}(\mathcal{V})$  for counting the numbers  $n_i$ .

**Proposition 3.3.** *If  $\Gamma_{f,g}(\mathcal{V})$  is empty, then  $n_i = 0$ , for all  $i \in \{1, \dots, q\}$ .*

**Proof.** Let  $V_i$  be a stratum of  $\mathcal{V}$  and  $x$  be a critical point of  $g|_{V_i \cap f^{-1}(\delta) \cap B_\epsilon}$ . Then, if  $\tilde{f}$  and  $\tilde{g}$  are analytic extensions of  $f$  and  $g$  in a neighborhood of  $x$ , respectively,  $x \in V_i \cap f^{-1}(\delta) \cap B_\epsilon$  and  $rk(d_x \tilde{f}|_{V_i}, d_x \tilde{g}|_{V_i}) \leq 1$ , that is,

$$x \in (V_i \cap f^{-1}(\delta) \cap B_\epsilon) \cap (\Sigma_{\mathcal{V}}f \cup \Sigma_{\mathcal{V}}g \cup \tilde{\Gamma}_{f,g}(\mathcal{V})).$$

By Proposition 1.3 of [12],  $\Sigma_{\mathcal{V}}f \subset \{f = 0\}$ . Therefore,

$$\Sigma g|_{V_i \cap f^{-1}(\delta) \cap B_\epsilon} = V_i \cap f^{-1}(\delta) \cap B_\epsilon \cap (\tilde{\Gamma}_{f,g}(V_i) \cup \Sigma g|_{V_i}) \subseteq V_i \cap f^{-1}(\delta) \cap B_\epsilon \cap \Gamma_{f,g}(V_i).$$

Since  $\Gamma_{f,g}(\mathcal{V})$  is empty, by Proposition 3.1,  $\Sigma g|_{V_i \cap f^{-1}(\delta) \cap B_\epsilon}$  is empty. Therefore,  $n_i = 0$ , for all  $i \in \{1, \dots, q\}$ .  $\square$

In [3], Dutertre and Grulha proved a Lê-Greuel type formula for the Brasselet number, with which it is possible count the number of stratified Morse critical points using Brasselet numbers. We apply their result to obtain a relation between Brasselet number in the setting we already know the number of Morse points. First let us show a more general result.

**Corollary 3.4.** *If  $\beta : X \rightarrow \mathbb{Z}$  is a constructible function with respect to the good stratification  $\mathcal{V}$  of  $X$  relative to  $f$  and  $\Gamma_{f,g}(\mathcal{V})$  is empty, then*

$$\chi(X \cap f^{-1}(\delta) \cap B_\epsilon, \beta) = \chi(X \cap \{g = 0\} \cap f^{-1}(\delta) \cap B_\epsilon, \beta).$$

**Proof.** By Corollary 3.2, since  $\Gamma_{f,g}(\mathcal{V})$  is empty,  $g$  is prepolar at the origin with respect to  $\mathcal{V}$  and, by Proposition 1.12 of [12], tractable at the origin with respect to  $\mathcal{V}$ . Then, by Theorem 4.2 of [3], we obtain

$$\chi(X \cap f^{-1}(\delta) \cap B_\epsilon, \beta) = \chi(X \cap \{g = 0\} \cap f^{-1}(\delta) \cap B_\epsilon, \beta) + \sum_{i=1}^q (-1)^{d-1} n_i \eta(V_i, \beta),$$

where  $n_i$  is the number of stratified Morse critical points of a Morsefication of  $g|_{V_i \cap f^{-1}(\delta) \cap B_\epsilon}$  appearing in  $V_i \cap f^{-1}(\delta) \cap B_\epsilon$ . Using again that  $\Gamma_{f,g}(\mathcal{V})$  is empty,  $n_i = 0$ , for all  $i \in \{1, \dots, q\}$  and we conclude the equality.  $\square$

If the constructible function  $\beta$  is giving by the local Euler obstruction, one obtains a relation between Brasselet numbers.

**Corollary 3.5.** *If  $X$  is equidimensional and  $\Gamma_{f,g}(\mathcal{V})$  is empty, then  $B_{f,X}(0) = B_{f,X^g}(0)$ .*

**Proof.** By Theorem 4.4 of [3],  $B_{f,X}(0) = B_{f,X^g}(0) + (-1)^{d-1} n_q$ , where  $n_q$  is the number of stratified Morse criticalpoints of a Morsefication of  $g|_{X \cap f^{-1}(\delta) \cap B_\epsilon}$  appearing on  $V_q \cap f^{-1}(\delta) \cap B_\epsilon$ . Since  $\Gamma_{f,g}(\mathcal{V})$  is empty,  $n_q$  is zero and the equality holds.  $\square$

When  $f$  is a generic linear form on  $\mathbb{C}^n$ ,  $B_{f,X}(0) = Eu_X(0)$  and  $B_{f,X^g}(0) = Eu_{X^g}(0)$ . Therefore, Corollary 3.5 implies the following consequence.

**Corollary 3.6.** *If  $X$  is equidimensional and  $\Gamma_{f,g}(\mathcal{V})$  is empty, then  $Eu_X(0) = Eu_{X^g}(0)$ .*

When both  $f$  and  $g$  has isolated singularity at the origin, Dutertre and Grulha proved several formulas about the Brasselet numbers of  $f$  and  $g$ . Using this formulas, one can obtain further information about these numbers if  $\Gamma_{f,g}(\mathcal{V})$  is empty. If that is the case, then  $g$  is prepolar at the origin with respect to the good stratification  $\mathcal{V}$  of  $X$  induced by  $f$ , given as a refinement of a Whitney stratification  $\mathcal{W} = \{W_i\}_i$  of  $X$ . By Corollary 6.1 of [3],  $f$  is prepolar at the origin with respect to the good stratification  $\overline{\mathcal{V}}$  induced by  $g$ , also given by a

refinement of  $\mathcal{W}$ . Applying Proposition 1.12 of [12], we obtain that  $\Gamma_{f,g}(\mathcal{V}) = \tilde{\Gamma}_{f,g}(\mathcal{V})$  and  $\Gamma_{g,f}(\overline{\mathcal{V}}) = \tilde{\Gamma}_{g,f}(\overline{\mathcal{V}})$ . But

$$\begin{aligned}
 \tilde{\Gamma}_{f,g}(\mathcal{V}) &= \bigcup_{V_i \in \mathcal{V}} \overline{\Sigma(f, g)|_{V_i \setminus (\{f=0\} \cup \{g=0\})}} \\
 &= \bigcup_{W_i \in \mathcal{W}} \overline{\Sigma(f, g)|_{(W_i \setminus \{f=0\}) \setminus \{g=0\}}} \\
 &= \bigcup_{W_i \in \mathcal{W}} \overline{\Sigma(f, g)|_{(W_i \setminus \{g=0\}) \setminus \{f=0\}}} \\
 &= \bigcup_{\overline{V}_i \in \overline{\mathcal{V}}} \overline{\Sigma(f, g)|_{\overline{V}_i \setminus (\{g=0\} \cup \{f=0\})}} \\
 &= \tilde{\Gamma}_{g,f}(\overline{\mathcal{V}}).
 \end{aligned}$$

Therefore, these four polar varieties are equal. Using this description, one concludes the following.

**Proposition 3.7.** *If  $\beta : X \rightarrow \mathbb{Z}$  is a constructible function with respect to the Whitney stratification  $\mathcal{W}$  and  $\Gamma_{f,g}(\mathcal{V})$  is empty, where  $\mathcal{V}$  is the good stratification of  $X$  induced by  $f$ , then*

$$\chi(X \cap f^{-1}(\delta) \cap B_\epsilon, \beta) = \chi(X \cap g^{-1}(\delta) \cap B_\epsilon, \beta).$$

**Proof.** Since  $\Gamma_{f,g}(\mathcal{V})$  is empty,  $g$  is prepolar at the origin with respect to  $\mathcal{V}$ . Then,  $f$  is prepolar at the origin with respect to the good stratification  $\overline{\mathcal{V}}$  induced by  $g$ . By Theorem 6.4 of [3],

$$\begin{aligned}
 \chi(X \cap f^{-1}(\delta) \cap B_\epsilon, \beta) &= \chi(X \cap g^{-1}(\delta) \cap B_\epsilon, \beta) \\
 &\quad + \sum_{i=1}^q (-1)^{d_i-1} (n_i - m_i) \eta(W_i, \beta),
 \end{aligned}$$

where  $d_i$  denotes the dimension of  $W_i \in \mathcal{W}$ . By Proposition 3.4, if  $m_i$  is the number of stratified Morse critical points of a Morsefication of  $f|_{X \cap g^{-1}(\delta) \cap B_\epsilon}$  appearing on  $V_i \cap g^{-1}(\delta) \cap \{f \neq 0\} \cap B_\epsilon$ , for each  $i \in \{1, \dots, q\}$ , since  $\Gamma_{f,g}(\mathcal{V})$  is empty,  $\Gamma_{f,g}(\overline{\mathcal{V}})$  is empty and  $m_i = 0$ . Since the number  $n_i$  of stratified Morse critical points of a Morsefication of  $g|_{X \cap f^{-1}(\delta) \cap B_\epsilon}$  in  $V_i \cap f^{-1}(\delta) \cap \{g \neq 0\} \cap B_\epsilon$  is zero, for each  $i \in \{1, \dots, q\}$ , the equality is proved.  $\square$

**Corollary 3.8.** *If  $X$  is equidimensional and  $\Gamma_{f,g}(\mathcal{V})$  is empty, then  $B_{g,X}(0) = B_{f,X}(0)$ .*

**Proof.** Since  $\Gamma_{f,g}(\mathcal{V})$  is empty,  $g$  (resp.  $f$ ) is prepolar with respect to the good stratification of  $X$  induced by  $f$  (resp.  $g$ ). Applying Corollary 6.5 of [3], we obtain  $B_{f,X}(0) = B_{g,X}(0) + (-1)^{d-1} (n_q - m_q)$ , where  $n_q$  (resp.  $m_q$ ) is the number of stratified Morse critical points of a Morsefication of  $g|_{X \cap f^{-1}(\delta) \cap B_\epsilon}$  (resp.  $f|_{X \cap g^{-1}(\delta) \cap B_\epsilon}$ ) appearing on the top stratum  $V_q \cap f^{-1}(\delta) \cap B_\epsilon$  (resp.  $V_q \cap g^{-1}(\delta) \cap B_\epsilon$ ). Using again that  $\Gamma_{f,g}(\mathcal{V})$  is empty, we have that  $n_q = m_q = 0$ , what leads to the equality.  $\square$

## Acknowledgements

The author thank the financial support from FAPESP-Brazil, Grants 2015/25191-9 and 2017/18543-1. This work was supported by the Research Institute for Mathematical Sciences, a Joint Usage/Research Center located in Kyoto University.

## References

- [1] BRASSELET, J.P., LÊ, D.T., SEADE, J., *Euler obstruction and indices of vector fields*, *Topology*, v.39, (2000), p. 1193-1208.
- [2] BRASSELET, J-P., MASSEY, D., PARAMESWARAN, A.J. AND SEADE, J. *Euler obstruction and defects of functions on singular varieties*, *Journal of the London Mathematical Society*, Cambridge University Press, 70, n.1, (2004), p. 59-76.
- [3] DUTERTRE, N., GRULHA JR., N.G., *Lê-Greuel type formula for the Euler obstruction and applications*, *Adv. Math.* 251 (2014), p. 127-146.
- [4] GORESKEY, M. AND MACPHERSON, R. *Stratified Morse theory*, Springer, v. 40, n. 1, (1988).
- [5] GREUEL, G-M. *Der Gauß-Manin-Zusammenhang isolierter Singularitäten von vollständigen Durchschnitten*, *Mathematische Annalen*, Springer, v. 214, n. 3, (1975), p. 235-266.
- [6] HAMM, H. *Lokale topologische Eigenschaften komplexer Räume*, *Mathematische Annalen*, v.191, n.3, (1971), p. 235-252.
- [7] LÊ, D.T. *Calcul Du Nombre de Milnor D'und Singularité Isolée D'intersection Complète*, *Centre de Mathematiques de l'Ecole Polytechnique*, (1973).
- [8] LÊ D. T., TEISSIER, B. *Variétés polaires locales et classes de Chern des variétés singulieres*, *Annals of Mathematics*, 114, No 3, (1981), p.457-491.
- [9] LOESER, F., *Formules intégrales pour certains invariants locaux des espaces analytiques complexes*, *Comment. Math. Helv.* 59(2)(1984) p. 204-225.
- [10] MACPHERSON, R. *Chern classes for singular algebraic varieties*, *Annals of Mathematics*, JSTOR, (1974), p. 423-432.
- [11] MASSEY, D. *Enriched Relative Polar Curves and Discriminants*, (2006).
- [12] MASSEY, D., *Hypercohomology of Milnor Fibers*, *Topology*, 35 (1996), no. 4 p. 969-1003.
- [13] MASSEY, D. *IPA-deformations of functions on affine space*, *Hokkaido Math. J.* 47 (2018), no. 3, 655-676.
- [14] MASSEY, D. *Numerical control over complex analytic singularities*, *American Mathematical Soc.*, (2003).
- [15] MILNOR, W. J. , *Singular Points of Complex Hypersurfaces*, *Annals of Mathematics Studies*, **25**, New Jersey, (1968).
- [16] MILNOR, W. J. , *Morse Theory - Based on lecture notes by M. Spivak and R. Wells*, *Annals of Mathematics Studies*, **51**, New Jersey, (1963).

- [17] SEADE, J., TIBĚR, M. AND VERJOVSKY, A. *Milnor numbers and Euler obstruction*, *Bulletin of the Brazilian Mathematical Society*, Springer, v. 36, n. 2, (2005), p. 275-283.
- [18] SCHÜRMANN, J. AND TIBAR, M. *Index formula for MacPherson cycles of affine algebraic varieties*, *BTohoku Mathematical Journal, Second Series*, Mathematical Institute, Tohoku University, v. 62, n. 1, (2010), p. 29-44.

Hellen Monção de Carvalho Santana

Institute of Mathematics and Computer Science - University of São Paulo. São Carlos. Brazil  
email: [hellenmcarvalho@hotmail.com](mailto:hellenmcarvalho@hotmail.com)